



BOTRYTIS (BUD ROT) | FUSARIUM | DOWNY MILDEW

**REDUCE DISEASE INCIDENCE & SEVERITY**

Use early for disease prevention & overall plant health. Apply 2-3 applications mid-flower to prevent *Botrytis* Bud Rot in late flower. Apply late in season to avoid issues of pesticide residues at harvest.

- ✓ Broad spectrum control
- ✓ Multiple modes of action
- ✓ Lipopeptides inhibit spore germination
- ✓ Harvest flexibility & worker friendly

**BASIC USE INSTRUCTIONS**

Foliar: 1-2 tbsp./1-2.5 gal. of water per 1,000 sq. ft.  
Soil Drench/Chemigation: 2 tbsp./gal. of water per 1,000 sq. ft.



**1-2 tbsp.**

or **15-30 ml.** per **gal.** per 1000 sq. ft.  
*Always read & follow label instructions.*

**GROWER RESOURCES:**



➔ **IPM Guides**

➔ **Educational Videos**

Scan QR codes with your camera app or go to [MarroneBio.com](http://MarroneBio.com)



FUNGICIDE | BACTERICIDE | ALGAECIDE | DISINFECTANT

**BOOST FUNGICIDE PERFORMANCE**

A broad spectrum peroxyacetic acid (PAA) suppresses, eliminates, & controls algae, fungi, & bacterial diseases. An excellent compliment to most fungal and bacterial disease IPM programs.

- ✓ Highly stable PAA
- ✓ Curative contact fungicide, bactericide & algaecide
- ✓ Kills pathogen spores on berry surface
- ✓ No residue
- ✓ Tank mix compatible\*

*Use caution when mixing with live organisms, as it can kill certain ones on contact. Jet-Ag 5% can be mixed with Stargus Biofungicide as long as it is sprayed right away.*



**Talk With An Expert**



**Brian Guess**

**949-316-9936**

**[bguess@profarmgroup.com](mailto:bguess@profarmgroup.com)**

# IPM GUIDE FOR INDOOR GROWING



FORMERLY MARRONE BIO INNOVATIONS

[ProFarmGroup.com](http://ProFarmGroup.com)

Learn more at [ProFarmGroup.com](http://ProFarmGroup.com)

©2023 Pro Farm Group Inc. All rights reserved. Grandevo, Stargus, Regalia & Venerate are registered trademarks of Marrone Bio Innovations, Inc. Agri-Mek SC® & Epi-Mek® are registered trademarks of Syngenta. Activator™ is a trademark of NuFarm. Exirel® is a registered trademark of FMC Agricultural Solutions. Verify to ensure the product you want to use is approved in your state.  
\*ONLY WHERE LEGAL. 02252023



POWDERY MILDEW | PYTHIUM ROT | BOTRYTIS

### FUNGICIDE & PLANT HEALTH

Use early for disease prevention & overall plant health & late in season to avoid issues of pesticide residues at harvest.

- ✓ Better yield & harvest quality
- ✓ Stimulates plant immunity & fights disease
- ✓ Quality control tested to ensure no human microbial pathogens
- ✓ Tank mix friendly: Can be mixed with Grandevo® CG Bioinsecticide or Venerate® CG Bioinsecticide

### BASIC USE INSTRUCTIONS

Transplant Dip: Roots & foliage benefit from a quick dip in a solution of 1 tbsp./gal.

Drench: 1-2 tbsp./gal. Every 14 days.

Foliar: 2 tbsp./gal. of water per 1,000 sq. ft. Apply at 7-day intervals.




**2 tbsp.**

or **30 ml.** per gal. • 1-2 gal. per 1000 sq. ft.  
*Always read & follow label instructions.*

### POWDERY MILDEW & BOTRYTIS BUD ROT - CANNABIS

Farm Forest Research



■ PM Incidence ■ Bud Rot Incidence  
Three applications made on a 7-day schedule in 42 gal./A



MITES | THRIPS | APHIDS

### MODERN PEST MANAGEMENT

- ✓ Prevents crop damage when used early
- ✓ Reduces reproduction, induces mortality, stops feeding & repels insects
- ✓ Soft on beneficials
- ✓ Tank mix friendly: Rotate with Venerate CG Bioinsecticide



### BASIC USE INSTRUCTIONS


Foliar: 2-4 tbsp./gal. of water per 1,000 sq. ft. Apply at 7-day intervals. Alternate with Venerate® CG Bioinsecticide.



**2-4 tbsp.**

or **28.6-57.2 grams** per gal. per 1000 sq. ft.  
*Always read & follow label instructions.*

### GRANDEVO® CG BIOINSECTICIDE & VENERATE® CG BIOINSECTICIDE ON MITES AVERAGE #/LEAFLET



MITES | THRIPS | APHIDS | ARMYWORMS

### IMRPOVE INSECT CONTROL

- ✓ Apply weekly to combat key pests
- ✓ Protects against sucking & chewing insects & mites
- ✓ Increases crop yield & quality
- ✓ Tank mix friendly: Rotate with Grandevo CG Bioinsecticide



### BASIC USE INSTRUCTIONS

Drench: 2.5-5 tbsp./gal. every 10 to 14-days.

Foliar: 2-5 tbsp./gal. of water per 1,000 sq. ft. Apply at 7-day intervals. Alternate with Grandevo CG Bioinsecticide.



**2.5-5 tbsp.**

or **37.5-75 ml.** per gal. per 1000 sq. ft.  
*Always read & follow label instructions.*

### VENERATE® CG BIOINSECTICIDE CG AGAINST THRIPS

Dr. Van Steenwyk, UC Berkeley

