



- For Vegetables, Shrubs, Lawns & Flowers
- Kills Tomato Hornworms, Bagworms,
   Armyworms, Webworms, Gypsy Moth Larvae, Cankerworms,
   Loopers, Tent Caterpillars & Other Listed Worms

# KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

# ACTIVE INGREDIENT

#### Home and Residential Use Only.

**Potency:** The percent active ingredient does not indicate product performance and potency rates are not federally standardized.



10859

**NET WEIGHT 4 LBS. (1.8 KG)** 



FIRST AID	
If in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul> <li>Take off contaminated clothing immediately.</li> <li>Rinse skin immediately with plenty of water for 15 -20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

In case of medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers or for product use information, call the National Pesticide Information Center at 1-877-757-4943. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency contact ChemTel Inc. ChemTel North America 1-800-255-3924



### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if inhaled. Avoid breathing dust. Wash thoroughly with soap and water after handing and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

## **ENVIRONMENTAL HAZARDS**

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

- 1. WHAT IS THIS PACKAGE FOR? Use ferti•lome® DIPEL® DUST Biological Insecticide to kill worms biologically. One bite of treated foliage and within minutes worms stop feeding and eventually die. ferti•lome@ **DIPEL® DUST Biological Insecticide** is easy to use for killing Tomato Hornworms, Bagworms, Armyworms, Webworms, Gypsy Moth Larvae, Cankerworms, Loopers, Tent Caterpillar, Tomato Fruitworms, Sod Webworms, Variegated Cutworms, Imported Cabbageworms, Rindworms, Melonworms and other listed worms.
- 2. HOW DO YOU USE THIS? Simply dust this product on plants at the rates listed below. Dust to thoroughly cover all plant surfaces. Apply when pests appear and repeat as necessary. Apply the specified number of ounces (ozs.) from the following chart up to 50 sq. ft. for vegetable and turf treatment

BEANS, PEAS, CABBAGE, TURNIPS, CAULIFLOWER, COLLARDS, BROCCOLI, AND MUSTARD GREENS: Use 1/2 oz. to kill Loopers and Webworms. For Hornworms, Cabbageworms, Diamondback Moths and Green Cloverworms, use 1/4 - 1/2 oz. For Armyworms, use 1-2 ozs.

TOMATOES, CUCUMBERS, SQUASH, PEPPERS, MELONS, LETTUCE, CELERY, POTATOES, AND SPINACH: Use 1/2 - 1 oz. to kill Loopers, Tomato Fruitworms, Variegated Cutworms, Rindworms, and Salt Marsh Caterpillars. For Hornworms, use 1/4 - 1/2 oz. For Armyworms and Melonworms, use 1-2 ozs.

TURF: To kill Sod Webworms use 2-4 ozs.

SHRUBS AND FLOWERS: Dust to thoroughly cover plant surfaces. Use to kill Loopers, Gypsy Moths, Cankerworms, Tent Caterpillars, Omnivorous Leafrollers, Tobacco Budworms, Spruce Budworms, Saddle Prominent Caterpillars, Fruit Tree Leafroller, Blackheaded Budworms, Saddleback Caterpillar, Western Tussock Moth, Diamondback Moth, Hornworms, California Oak Moths and Armyworms.

3. HERE ARE THE RESULTS OF WHAT YOU EXPECT! When used as directed , ferti•lome® DIPEL® DUST Biological Insecticide will kill a wide variety of leaf-eating caterpillars that can severely damage and kill vegetables, shrubs, lawn and flowers.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. PESTICIDE STORAGE: Store in a cool and dry place inaccessible to children and pets

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418 EPA Reg. No.7401-290 • EPA Est. No. 7401-TX-01 Visit Us At: www.fertilome.com Product Questions? 855-270-4776

10859-0815-IC

